

175 Cambridge Street Boston, MA 02114 617-643-9999 www.mghsportsmedicine.org

Shoulder Rotator Cuff and Scapular Strengthening Program

External Rotation

Attach the theraband at waist level to a doorknob or post. While standing sideways to the door and facing straight ahead, grasp one end of the band and pull the band all the way through until it is taut. Feet are shoulder width apart and the knees are slightly flexed. The elbow is placed next to the side with the hand as close to your chest as possible (think of this elbow as being a hinge on a gate). Taking the cord in the hand 'set' the shoulder blade and move the hand away from the body as far as it feels comfortable. Return to the start position.

Internal Rotation

Attach the Theraband at waist level in a doorknob or post. While standing sideways to the door and looking straight ahead, grasp one end of the handle and pull the cord all the way through until it is taut. Feet are shoulder width apart and the knees are slightly flexed. The elbow is placed next to the side and is flexed at 90 degrees (think of this elbow as being a hinge on a gate). Taking the cord in the hand, 'set' the shoulder blade and move the hand toward the belly as far as it feels comfortable, or to where the endpoint of pain limits you. Return to the start position.









Lateral Raises

Stand with the arm at your side with the elbow straight and the hands rotated so that the thumbs face forward. Raise the arm straight out to the side, palm down, until the hands reach shoulder level. Do not raise the hands higher than the shoulder. Pause and slowly lower the arm.

<u>Standing forward flexion ('full-can')</u> exercise

Stand facing a mirror with the hands rotated so that the thumbs face forward. While keeping the shoulder blade 'set' and keeping the elbows straight, raise the arms forward and upward to shoulder level with a slight outward angle (30°). Pause for one second and slowly lower and repeat.









GENERAL HOSPITAL
Sports Physical Therapy

MASSACHUSETTS

175 Cambridge Street Boston, MA 02114 617-643-9999 www.mghsportsmedicine.org

Sidelying external rotation

Lying on the non-operated side, bend your elbow to a 90-degree angle and keep the operated arm firmly against your side with your hand resting on your abdomen. By rotation at the shoulder, raise your hand upward, toward the ceiling through a comfortable range of motion. Hold this position for 1 to 2 seconds, and then slowly lower the hand.





Prone extension

The starting position for this exercise is to bend over at the waist so that the affected arm is hanging freely straight down. Alternatively, lie face down on your bed with the operated arm hanging freely off of the side. While keeping the shoulder blade 'set' and keeping the elbow straight, raise the arm backward toward your hip with the thumb pointing outward. Do not lift your hand past the level of your hip.



Prone rowing exercise

The starting position for this exercise is to bend over at the waist so that the affected arm is hanging freely straight down. Alternatively, lie face down on your bed with the operated arm hanging freely off of the side. While keeping the shoulder blade 'set', raise the arm up toward the ceiling while bending at the elbow. The elbow should be drawn along the side of the body until the hands touch the lower ribs. Always return slowly to the start position.







175 Cambridge Street Boston, MA 02114 617-643-9999 www.mghsportsmedicine.org

Prone horizontal abduction ('T's)

The starting position for this exercise is to bend over at the waist so that the affected arm is hanging freely straight down. Alternatively, lie face down on your bed with the operated arm hanging freely off of the side. Rotate your hand so that the thumb faces forward. While keeping the shoulder blade 'set' and keeping the elbows straight, slowly raise your arm away from your body to shoulder height, through a pain-free range of motion (so that your hand now has the thumb facing forward, and aligned with your cheek .) Hold that position for 1 to 2 seconds and slowly lower. Limit the height that you raise the arm to 90 degrees, or in other words, horizontal to the floor.




Prone scaption ('Y's)

The starting position for this exercise is to bend over at the waist so that the affected arm is hanging freely straight down. Alternatively, lie face down on your bed with the operated arm hanging freely off of the side. Keep the shoulder blade 'set' and keep the elbows straight. Slowly raise the arm away from your body and slightly forward through a pain-free range of motion (so that your hand now has the thumb facing up, and is aligned with your forehead). Hold that position for 1 to 2 seconds and slowly lower. Limit the height that you raise the arm to 90 degrees, or in other words, horizontal to the floor.

Prone external rotation at 90° abduction ('U's)

Lie face down on a table with your arm hanging over the side of the table. Raise the arm to shoulder height at a 90° angle to the body. While holding the arm in this position, rotate the hand upward, until the hand is even with the elbow. Hold one second and slowly let the hand rotate to the starting position and repeat.










MASSACHUSETTS GENERAL HOSPITAL Sports Physical Therapy 175 Cambridge Street Boston, MA 02114 617-643-9999 www.mghsportsmedicine.org

'W's

With the tubing attached in front of you, stand with the tubing in both hands with the elbows bent at 90° and fixed at your side. Pull the band outward, keeping the elbow at your side. The arms rotate outward making the shape of a 'W'.



Dynamic Hug

With the tubing attach behind you at shoulder height, grip both ends of the tubing in your hands with the tubing on the outside of your shoulders. Pull the band forward and slightly downward in a 'hugging' motion, or as if you were wrapping both arm around a small tree. Pause and return slowly to the starting position.



