

A Service of the Massachusetts Smokers' Helpline

In Collaboration with the Massachusetts Department of Public Health and Massachusetts Health Plans

- If a patient is interested in quitting smoking, fill out this form with them and Fax/Send to 1-866-560-9113.
- The Massachusetts Smokers' Helpline offers free cessation services (counseling and NRT) and sends feedback reports to the provider as noted below.
- Inform patients that QuitWorks calls will come from 617-262-2200.
 Four attempts will be made to reach the patient.
- This program is free for all Massachusetts residents regardless of insurance status



Massachusetts Referral Form

	lame			Are you 18 ☐ Yes ☐ N or older?	
Mailing Address	City		State	Zip	
()					
Phone Number					
When should we call? (check all that apply)	☐ Morning	☐ Afternoon	Evening	☐ No pı	reference
Language Preference:	anish 🔲 O	ther (specify)			
May we leave a message? ☐ Yes ☐ No					
Primary Insurance		ufts Health Plan	Health Plan 🔲 Harvard Pilgrim		
		ther	■ None	□ None	
QuitWorks program. I also authorize QuitWorks to di care provider listed on this form. Patient Signature	sciose information	i about my progre	ss in attempting t	o quit smo	king to the health
Health Care Providers: Complete this:	section				
Referring Provider:			Phone Number ()		
Facility:			Fax Numb	or	
Address:			()		
Send feedback report to:					
─ Same as		()		()	1
above orName		Phone Number		Fax Nur	nher
PEDIATRICS ONLY:				· ux itui	
Patient's relationship to child:	☐ Father ☐	Other (specify)			
Child/Children's name: (to help with records					

Copies of this form can be downloaded from WWW.QUITWORKS.ORG



Quick Guide To Pharmacotherapy In Tobacco Treatment

NICOTINE REPLACEMENT OPTIONS

Combining long-acting NRT (patch) with a short-acting NRT (gum, lozenge, or inhaler) is more effective than using a single type of NRT

LONG-ACTING PRODUCTS

PATCHES

21 mg, 14 mg, 7 mg Dose: 1 patch every 24 hrs. Duration: Start: 21 mg patch if ≥10 cig/day 6-14 wks

14 mg patch if < 10 cig/day

SHORT-ACTING PRODUCTS

GUM

2mg, 4 mg 1 piece every 1-2 hrs. Duration: Dose:

24 pieces/day 6-14 wks Max:

LOZENGE or MINI-LOZENGE

1 lozenge every 1-2 hrs. 2mg, 4 mg Dose: Duration:

Max: 20 pieces/day 12 wks

NASAL SPRAY (Nicotrol® NS)

10 mg/ml 1-2 doses per hr. Duration: Dose:

Max: 5 doses/hr or 40 doses/day 3-6 mos

INHALER (Nicotrol® Inhaler)

6-16 cartridges/day Duration: Dose: 16 cartridges/day Max: 3-6 mos

BUPROPION SR

(Zvban®/ Wellbutrin SR®)

May be combined with nicotine replacement

150 mg tablets Dose: 150 mg once per day (days 1-3) Duration: 12 wks*

150 mg twice per day (day 4+)

300 mg/day Max:

VARENICLINE (Chantix®)

0.5 mg, 1 mg tablets Dose: Starting Month Pak = Duration:

> 0.5 mg once per day (days 1-3) 12 wks*

0.5 mg twice per day (days 4-7) 1 mg twice per day (days 8+)

Continuing Month Pak = 1 mg twice per day

Max: 2 mg/day

Source: Fiore MC, Jaen CR, Baker TB, et al. Treating Tobacco Use and Dependence: 2008 Update. Quick Reference Guide for Clinicians, Rockville, MD: U.S. Department of Health and Human Services. Public Health Service. April 2009. Inclusion of this adult dosage chart is strictly for the convenience of the prescribing provider. Please consult the Physicians' Desk Reference for complete product information and contraindications. This chart does not indicate or authorize insurance benefit coverage for any of these medications. This chart is provided by the Massachusetts Department of Public Health's Tobacco Cessation and Prevention Program. For insurance benefit information, the patient will need to contact his/her insurer directly. The cost or provision of these medications is not included as any part of the Massachusetts Smokers' Helpline or QuitWorks program.

^{*} If quit at 12 wks, consider 12 more weeks of drug