

## Rehabilitation Protocol for SLAP Repair-Type II

This protocol is intended to guide clinicians through the post-operative course for SLAP Repair-Type II. This protocol is time based (dependent on tissue healing) as well as criterion based. Specific intervention should be based on the needs of the individual and should consider exam findings and clinical decision making. The timeframes for expected outcomes contained within this guideline may vary based on surgeon's preference, additional procedures performed, and/or complications. If a clinician requires assistance in the progression of a post-operative patient, they should consult with the referring surgeon.

The interventions included within this protocol are not intended to be an inclusive list. Therapeutic interventions should be included and modified based on the progress of the patient and under the discretion of the clinician.

### Considerations for the Post-operative SLAP repair Rehabilitation Program

Many different factors influence the post-operative SLAP repair rehabilitation outcome, including the type of SLAP lesion, the size of the tear/number of anchors placed, concomitant procedures and amount of shoulder hypermobility and/or hyperlaxity. Consider mechanism of injury, as well as the sport the athlete would like to return to when initiating certain interventions.

### Post-operative Complications

If you develop a fever, unresolving numbness/tingling, excessive drainage from the incision, uncontrolled pain or any other symptoms you have concerns about you should contact the referring physician.

### PHASE I: IMMEDIATE POST-OP (0-3 WEEKS AFTER SURGERY)

<b>Rehabilitation Goals</b>	<ul style="list-style-type: none"> <li>• Protect surgical repair</li> <li>• Reduce swelling, minimize pain</li> <li>• Maintain UE ROM in elbow, hand and wrist</li> <li>• Gradually increase shoulder PROM</li> <li>• Minimize muscle inhibition</li> <li>• Patient education</li> </ul>
<b>Sling</b>	<ul style="list-style-type: none"> <li>• Neutral rotation</li> <li>• Use of abduction pillow in 30-45 degrees abduction</li> <li>• Use at night while sleeping</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>• No shoulder AROM/AAROM</li> <li>• No elbow AROM (avoid biceps contraction)</li> <li>• No lifting of objects</li> <li>• No supporting of body weight with hands</li> <li>• No reaching behind back</li> </ul>
<b>Interventions</b>	<p><i>Swelling Management</i></p> <ul style="list-style-type: none"> <li>• Ice, compression</li> </ul> <p><i>Range of motion/Mobility</i></p> <ul style="list-style-type: none"> <li>• PROM: ER&lt;30 scapular plane, forward elevation &lt;90, full elbow flex and ext, <a href="#">seated GH flexion table slide</a>, <a href="#">horizontal table slide</a></li> <li>• AROM: hand, wrist</li> <li>• AAROM: none</li> </ul>

	<p><i>Strengthening (Week 2)</i></p> <ul style="list-style-type: none"> <li>Periscapular: <a href="#">scap retraction*</a>, <a href="#">prone scapular retraction*</a>, <a href="#">standing scapular setting*</a>, <a href="#">supported scapular setting</a>, <a href="#">inferior glide</a>, <a href="#">low row</a> <ul style="list-style-type: none"> <li>*to neutral; avoid shoulder extension</li> </ul> </li> <li>Rotator cuff: submaximal pain-free <a href="#">internal/external rotation isometrics</a></li> <li><a href="#">Ball squeeze</a></li> </ul>
<b>Criteria to Progress</b>	<ul style="list-style-type: none"> <li>90 degrees shoulder PROM forward elevation</li> <li>30 degrees of shoulder PROM ER in the scapular plane</li> <li>Full elbow PROM flexion and extension</li> <li>Palpable muscle contraction felt in scapular and shoulder musculature</li> <li>No complications with Phase I</li> </ul>

### **PHASE II: INTERMEDIATE POST-OP (4-6 WEEKS AFTER SURGERY)**

<b>Rehabilitation Goals</b>	<ul style="list-style-type: none"> <li>Continue to protect surgical repair</li> <li>Reduce swelling, minimize pain</li> <li>Gradually increase shoulder PROM</li> <li>Minimize substitution patterns with shoulder AAROM</li> <li>Initiate motor control exercise</li> <li>Patient education</li> </ul>
<b>Sling</b>	<ul style="list-style-type: none"> <li>Neutral rotation</li> <li>Use of abduction pillow in 30-45 degrees abduction</li> <li>Use at night while sleeping</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>No shoulder AROM</li> <li>No elbow AROM (avoid biceps contraction)</li> <li>No lifting of objects</li> <li>No supporting of body weight with hands</li> <li>No reaching behind back</li> </ul>
<b>Interventions</b> <i>*Continue with Phase I interventions</i>	<p><i>Range of motion/Mobility</i></p> <ul style="list-style-type: none"> <li>PROM: ER&lt;45 scapular plane, Forward elevation &lt;120, IR in scapular plane</li> <li>AAROM: <a href="#">Active assistive shoulder flexion</a>, <a href="#">shoulder flexion with cane</a>, <a href="#">cane external rotation stretch</a>, <a href="#">washcloth press</a></li> </ul> <p><i>Strengthening</i></p> <ul style="list-style-type: none"> <li>Periscapular: <a href="#">Row on physioball*</a>, <a href="#">shoulder extension on physioball*</a>, <a href="#">serratus punches</a> <ul style="list-style-type: none"> <li>*to neutral; avoid shoulder extension</li> </ul> </li> </ul> <p><i>Motor Control</i></p> <ul style="list-style-type: none"> <li>Rhythmic Stabilization: Internal and external rotation in scaption and forward elevation 90-120</li> </ul> <p><i>Stretching</i></p> <ul style="list-style-type: none"> <li><a href="#">Sidelying horizontal ADD</a>, <a href="#">sleeper stretch</a></li> </ul>
<b>Criteria to Progress</b>	<ul style="list-style-type: none"> <li>120 degrees shoulder PROM forward elevation</li> <li>45 degrees shoulder PROM ER in scapular plane</li> <li>Minimal substitution patterns with shoulder AAROM</li> <li>Pain &lt; 4/10</li> <li>No complications with Phase II</li> </ul>

### **PHASE III: INTERMEDIATE POST-OP CONT'd (7-8 WEEKS AFTER SURGERY)**

<b>Rehabilitation Goals</b>	<ul style="list-style-type: none"> <li>Do not overstress healing tissue</li> <li>Reduce swelling, minimize pain</li> <li>Gradually increase shoulder PROM/AAROM</li> <li>Initiate shoulder and elbow AROM</li> <li>Initiate RTC strengthening</li> <li>Improve scapular muscle activation</li> <li>Patient education</li> </ul>
-----------------------------	---

<b>Sling</b>	<ul style="list-style-type: none"> <li>Discontinue</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>No resisted elbow flexion</li> <li>No lifting of heavy objects (&gt;10 lbs)</li> </ul>
<b>Interventions</b> <i>*Continue with Phase I-II interventions</i>	<p><i>Range of motion/Mobility</i></p> <ul style="list-style-type: none"> <li>PROM: ER Full in scapular plane, ≤90 degrees ER in 90 degrees of abduction, IR Full in scapular plane/90 degrees of abduction, Forward elevation Full <ul style="list-style-type: none"> <li>*do not push beyond 90 degrees ER in 90 degrees of abduction</li> </ul> </li> <li>AAROM: <a href="#">seated shoulder elevation with cane</a>, <a href="#">seated incline table slides</a>, <a href="#">ball roll on wall</a></li> <li>AROM: <a href="#">supine flexion</a>, <a href="#">salutes</a>, <a href="#">supine punch</a>, <a href="#">wall climbs</a>, <a href="#">elbow flexion</a></li> </ul> <p><i>Strengthening</i></p> <ul style="list-style-type: none"> <li>Rotator cuff: <a href="#">side-lying external rotation</a>, <a href="#">standing external rotation w/ resistance band</a>, <a href="#">standing internal rotation w/ resistance band</a></li> <li>Periscapular: <a href="#">Resistance band shoulder extension*</a>, <a href="#">resistance band seated rows*</a>, <a href="#">lawn mowers</a>, <a href="#">robbery</a> <ul style="list-style-type: none"> <li>*to neutral; avoid shoulder extension</li> </ul> </li> <li>Elbow: <a href="#">Resistance band elbow extension</a></li> </ul> <p><i>Motor Control</i></p> <ul style="list-style-type: none"> <li><a href="#">Quadruped alternating isometrics</a></li> </ul>
<b>Criteria to Progress</b>	<ul style="list-style-type: none"> <li>Full pain-free shoulder PROM ER and forward elevation</li> <li>Within 10 degrees of shoulder IR PROM of contralateral shoulder</li> <li>Minimal substitution patterns with shoulder AROM</li> <li>Pain &lt; 4/10</li> </ul>

#### **PHASE IV: TRANSITIONAL POST-OP (9-12 WEEKS AFTER SURGERY)**

<b>Rehabilitation Goals</b>	<ul style="list-style-type: none"> <li>Do not overstress healing tissue</li> <li>Restore full shoulder PROM and AROM</li> <li>Initiate resisted elbow flexion at 12 weeks</li> <li>Improve dynamic shoulder stability</li> <li>Progress periscapular strength</li> <li>Gradually return to full functional activities</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>No lifting of heavy objects (&gt; 10 lbs)</li> </ul>
<b>Interventions</b> <i>*Continue with Phase II-III interventions</i>	<p><i>Range of motion/mobility</i></p> <ul style="list-style-type: none"> <li>PROM: Full</li> <li>AROM: <a href="#">Supine forward elevation with elastic resistance to 90 deg</a>, <a href="#">seated scaption</a> and <a href="#">seated flexion to 90 degrees elevation</a></li> </ul> <p><i>Strengthening</i></p> <ul style="list-style-type: none"> <li>Periscapular: <a href="#">Push-up plus on knees</a>, <a href="#">prone shoulder extension Is*</a>, <a href="#">resistance band forward punch</a>, <a href="#">tripod</a> <ul style="list-style-type: none"> <li>*to neutral; avoid shoulder extension</li> </ul> </li> <li>Elbow (12 weeks): <a href="#">resistance band bicep curls</a></li> </ul> <p><i>Motor control</i></p> <ul style="list-style-type: none"> <li>Ball stabilization on wall</li> </ul> <p><i>Stretching</i></p> <ul style="list-style-type: none"> <li><a href="#">Hands behind head</a>, <a href="#">IR behind back with towel</a>, <a href="#">triceps</a>, lats, <a href="#">doorway series</a></li> </ul>
<b>Criteria to Progress</b>	<ul style="list-style-type: none"> <li>Full pain-free shoulder PROM and AROM</li> <li>Minimal to no substitution patterns with shoulder AROM</li> <li>Performs all exercises demonstrating symmetric scapular mechanics</li> <li>Pain &lt; 2/10</li> </ul>

## PHASE V: LATE POST-OP (13-16 WEEKS AFTER SURGERY)

<b>Rehabilitation Goals</b>	<ul style="list-style-type: none"> <li>Maintain pain-free shoulder ROM</li> <li>Enhance functional use of upper extremity</li> </ul>
<b>Intervention</b> <i>*Continue with Phase II-IV interventions</i>	<p><i>Strengthening</i></p> <ul style="list-style-type: none"> <li>Rotator cuff: <a href="#">resistance band standing external rotation at 90 degrees</a>, <a href="#">resistance band standing internal rotation at 90 degrees</a></li> <li>Periscapular: <a href="#">T and Y</a>, <a href="#">push-up plus knees extended</a>, <a href="#">pointer/birddog</a>, <a href="#">wall push up</a>, <a href="#">resistance band Ws</a>, <a href="#">resistance band dynamic hug</a></li> </ul> <p><i>Motor Control</i></p> <ul style="list-style-type: none"> <li><a href="#">PNF – D1 diagonal lifts</a>, <a href="#">PNF – D2 diagonal lifts</a>, <a href="#">field goals</a>, <a href="#">resistance band PNF pattern</a>, <a href="#">PNF – D1 diagonal lifts w/ resistance</a>, <a href="#">wall slides w/ resistance band</a></li> </ul>
<b>Criteria to Progress</b>	<ul style="list-style-type: none"> <li>Clearance from MD and ALL milestone criteria below have been met</li> <li>Full pain-free shoulder PROM and AROM</li> <li>ER/IR strength minimum 85% of the uninvolved arm</li> <li>ER/IR ratio 60% or higher</li> <li>Negative impingement and instability signs</li> <li>Performs all exercises demonstrating symmetric scapular mechanics</li> <li>QuickDASH</li> <li>PENN</li> </ul>

## PHASE VI: EARLY RETURN-TO-SPORT (4-6 MONTHS AFTER SURGERY)

<b>Rehabilitation Goals</b>	<ul style="list-style-type: none"> <li>Maintain pain-free ROM</li> <li>Continue strengthening and motor control exercises</li> <li>Enhance functional use of upper extremity</li> <li>Gradual return to strenuous work/sport activity</li> </ul>
<b>Interventions</b> <i>*Continue with Phase II-V interventions</i>	<p><i>Strengthening</i></p> <ul style="list-style-type: none"> <li>See specific return-to-sport/throwing program (coordinate with physician)</li> </ul>
<b>Criteria to Progress</b>	<ul style="list-style-type: none"> <li>Last stage-no additional criteria</li> </ul>
<b>Return-to-Sport</b>	<ul style="list-style-type: none"> <li>For the recreational or competitive athlete, return-to-sport decision making should be individualized and based upon factors including level of demand on the upper extremity, contact vs non-contact sport, frequency of participation, etc. We encourage close discussion with the referring surgeon prior to advancing to a return-to-sport rehabilitation program.</li> </ul>

Revised 10/2021

<b>Contact</b>	Please email <a href="mailto:MGHSportsPhysicalTherapy@partners.org">MGHSportsPhysicalTherapy@partners.org</a> with questions specific to this protocol
----------------	--

### References:

- Ahsan, ZS, Hsu JE, et al. The Snyder classification of superior labrum anterior and posterior (SLAP) lesions. *Clin Orthop Relat Res*. 2016. 474:2075-2078.
- Christopher ZR, Kennedy J, et al. Rehabilitation and return to play following superior labral anterior to posterior repair. *Oper Tech Sports Med*. 2017. 25:132-144.
- Cools AM, Borms D, Cottens S, et al. Rehabilitation exercises for athletes with biceps disorders and SLAP lesions. *AJSM*. 2014. 42(6): 1315-1323.
- Itoi E, Hatakeyama Y, et al. Position of immobilization after dislocation of the shoulder. *JBJS*. 1999. 81-A(3): 385-390.
- Kibler, W.B., Sciascia, A. D., Uhl, T. L., et al. Electromyographic analysis of specific exercises for scapular control in early phases of shoulder rehabilitation. *The American Journal of Sports Medicine*. 2008. 36(9): p. 1789-1798.
- Wilk KE, Reinold MM, et al. Current concepts in the recognition and treatment of superior labral (SLAP) lesions. *JOSPT*. 2005. 35(5): 273-291.