


Daniel A. Haber, MD, PhD



Haber Laboratory*

Risa Burr, PhD
Dante Che
Charles Dai, MD
Douglas Fox, PhD
Sarah Guay
Hongshan Guo, PhD
Daniel Haber, MD, PhD
Elad Horwitz, PhD
Shih-Bo Huang, PhD
Justin Kelly
Laura Libby
Doug Micalizzi, MD
Ajinkya Revandkar, PhD
Hunter Russell
Karsen Seeger
Joanna Vuille**
Ben Wittner, PhD

* Co-directed with Shyamala Maheswaran, PhD

** Graduate student


The Haber laboratory focuses on understanding mutations that are acquired by tumors and render them susceptible to specific targeted drug therapies. In 2004, we identified mutations in the EGFR gene in lung cancers which confer dramatic sensitivity to drugs that specifically inhibit that pathway. This finding triggered the application of targeted therapies in lung cancer, and more generally pointed to the critical importance of mutational analysis for treatment selection in common epithelial cancers. Since then, we have collaborated with the bioengineering team led by Dr. Mehmet Toner, the molecular biology group of Dr. Shyamala Maheswaran, and the MGH Cancer Center clinical disease centers to develop, characterize and apply microfluidic devices to isolate rare circulating tumor cells (CTCs) in the blood of patients with cancer. Using these technologies, our lab seeks to explore 1) blood-based early detection of cancer, 2) noninvasive monitoring of cancer for the emergence of drug resistance, and 3) understanding mechanisms of tumor cell dissemination and metastasis, with the ultimate goal of suppressing blood-borne spread of cancer.

Our laboratory is interested in the genetics of human cancer. Current projects include the use of a microfluidic device to capture circulating tumor cells (CTCs) and its application in early detection of invasive cancer, molecular-directed therapy, and in the study of human cancer metastasis.

Circulating Tumor Cells and Molecular Genetics Underlying Targeted Cancer Therapeutics

Activating mutations in the epidermal growth factor receptor (*EGFR*) were identified in our laboratory in the subset of non-small cell lung cancer (NSCLC) with dramatic responses to the tyrosine kinase inhibitor gefitinib. We have studied mechanisms underlying such oncogene addiction, as well as the pathways that lead to the acquisition of resistance to targeted therapies, including the application of irreversible kinase inhibitors to circumvent mutations that alter drug binding affinity. Following these efforts to monitor the emergence of drug resistance

mutations, we established collaborations with the Toner and Maheswaran laboratories to characterize novel microfluidic devices capable of isolating CTCs from the blood of cancer patients. Our most advanced version of these CTC-Chips relies upon blood flow through a specialized chamber, which allows the high efficiency depletion of antibody-tagged leukocytes, thereby enriching for intact CTCs without selection bias. We have shown that the number of captured CTCs correlates with clinical evidence of tumor response, and that the cells can be used to define molecular markers characteristic of the underlying malignancy, including *EGFR* mutations in lung cancer and measurements of androgen receptor (AR) activity in prostate cancer. We have applied next generation single-molecule RNA sequencing and RNA-in-situ hybridization to characterize the heterogeneous expression profiles of individual CTCs in breast, prostate and pancreatic cancers, as well as melanoma and glioblastoma. To facilitate CTC



Circulating prostate tumor cell cluster stained for PSA (green) along with Ki67 (orange) and CD45 (red).

quantitation and provide the sensitivity and specificity required for early cancer detection, we have established a droplet digital PCR readout for CTC-derived RNA, with promising applications in the early detection of liver cancer.

In addition to noninvasive detecting and monitoring of cancer, CTCs provide a window to study the process of blood-borne metastasis. We demonstrated treatment-associated epithelial-to-mesenchymal transitions (EMT) within CTCs from women with breast cancer. Using a combination of mouse models and patient-derived studies, we observed that tumor-derived fragments generate CTC-Clusters, which have greatly enhanced metastatic propensity compared with single CTCs. CTC-Clusters are held together by plakoglobin, whose knockdown dramatically suppresses CTC-Cluster formation and metastatic spread of breast cancer cells. We successfully established long-term *in vitro* cultures of CTCs from patients with estrogen-receptor (ER)-positive breast cancer, identifying treatment-associated mutations in the

estrogen receptor (ESR1), as well as acquired mutations in druggable therapeutic targets, such as *PIK3CA* and *FGFR*. The development of such CTC-derived cultures may enable functional predictive drug testing, combined with detailed genetic analysis of tumor cells sampled noninvasively during the course of cancer treatment. In a recent study of metastasis initiation, we identified a critical role for the nuclear orphan receptor *NR4A1*, as a regulator of transcriptional elongation for replication stress-induced Immediate Early Genes (IEG). Both genetic and drug treatment studies point to *NR4A1* as a “non-oncogene dependent” target in tumorigenesis. Ongoing studies are directed at using patient-derived CTCs and mouse models to understand key steps in cancer metastasis, including the shift from cell quiescence to proliferation, viability during blood-borne transit, and resistance to targeted and immune therapies.

Selected Publications:

Guo H, Golczer G, Wittner BS, Langenbacher A,...Vasudevan S, Zou L, Mostoslavsky R, Maheswaran S*, Lawrence MS*, Haber DA*. NR4A1 regulates expression of immediate early genes, suppressing replication stress in cancer. *Mol Cell*. 81(19): 4041-4058, 2021 PMID: 34624217.

Hong X, Roh W, Sullivan RJ, Wong KHK, Wittner BS,...Toner M, Stott SL, Getz G, Maheswaran S*, Haber DA*. The lipogenic regulator SREBP2 induces transferrin in circulating melanoma cells and suppresses ferroptosis. *Cancer Discovery*. 11:678-95, 2021 PMID 33203734.

Ebright RY, Lee S, Wittner BS, Niederhoffer KL,...Ting DT, Toner M, Vasudevan S, Haber DA*, Maheswaran S, Micalizzi DS. Deregulation of ribosomal protein expression and translation promotes breast cancer metastasis. *Science*. 367(6485):1468-1473, 2020. PMID 32029688.

Zheng Y, Comaills V, Burr R, Boulay G, Miyamoto DT, Wittner BS,...Rivera MN, Maheswaran S, Haber DA. COX-2 mediates tumor-stromal prolactin signaling to initiate tumorigenesis. *Proc Natl Acad Sci USA*. 116:5223-5232, 2019. PMID: 30819896.

Miyamoto DT, Lee RJ, Kalinich M, LiCausi JA, Zheng Y,...Toner M, Maheswaran S, Haber DA. An RNA-based digital circulating tumor cell signature is predictive of drug response and early dissemination in prostate cancer. *Cancer Discovery*. 8: 288-303, 2018. PMID: 29301747.

Jordan NV, Bardia A, Wittner BS, Benes C, Ligorio M, Zheng Y, Yu M, Sundaresan TK, Licausi JA, Desai R, O’Keefe RM, Ebright RY, Boukhali M, Sil S, Onozato ML, Iafrate AJ, Kapur R, Sgroi D, Ting DT, Toner M, Ramaswamy S, Haas W, Maheswaran S, Haber DA. HER2 expression identifies dynamic functional states within circulating breast cancer cells. *Nature*. 537(7618):102-106, 2016.

*Co-corresponding authors