

Pediatric and Adolescent Rehabilitation Protocol for Post-Operative Management of Hip Labrum Repair for FAI Osteochondroplasty

This protocol is intended to guide clinicians through the post-operative course for the post-operative management of hip labrum repair for femoroacetabular impingement (FAI) and osteochondroplasty. This protocol is time based (dependent on tissue healing) as well as criterion based. Specific intervention should be based on the needs of the individual and should consider exam findings and clinical decision making. The timeframes for expected outcomes contained within this guideline may vary based on surgeon's preference, additional procedures performed, and/or complications. If a clinician requires assistance in the progression of a post-operative patient, they should consult with the referring surgeon.

The interventions included within this protocol are not intended to be an inclusive list of exercises. Therapeutic interventions should be included and modified based on the progress of the patient and under the discretion of the clinician.

Post-Operative Considerations:

Many different factors influence the post-operative management of hip labrum repair for FAI rehabilitation outcomes, including if a labrum was repaired or reconstructed. If the labral tissue is inadequate the surgeon may reconstruct the labrum using an autograft or allograft. This information can be accessed in the operative note and will impact rehabilitation. It is recommended that clinicians collaborate closely with the referring physician. One surgical technique that merits special consideration in post-operative rehabilitation is capsular closure. Capsular closure is performed to restore the normal anatomy and minimize the risk of postoperative issues with instability. With the capsular repair closure technique, it is necessary to protect and limit hip external rotation and extension in the early healing phase to protect the integrity of the repair. Capsular integrity has been correlated to improved outcomes after hip arthroscopy with FAI correction.

Complications that frequent the pediatric and adolescent population can include pudendal nerve palsy, and recurrent labral tears. It is also important to consider the implications of the anatomy and physiology of the adolescent athlete population in the pathogenesis and management of conditions such as FAI. There is higher prevalence of cam deformities in adolescents participating in sports compared with nonathlete cohorts. Additionally, as the pediatric and adolescent population is skeletally immature which the open apophyses are inherently weak, placing them at higher risk for avulsion injury and repair. Furthermore, any amount of resection of the anterolateral head-neck junction also increases the risk for femoral neck fracture, thus this should be monitored well when adhering to protocol post-operatively.

If the patient develops a fever, unresolving numbness/tingling, excessive drainage from the incision, uncontrolled pain or any other symptoms you have concerns about, the referring physician should be contacted.

Procedures Performed:

- | | |
|--|---|
| <input type="checkbox"/> Acetabuloplasty | <input type="checkbox"/> Chondroplasty |
| <input type="checkbox"/> Labral repair | <input type="checkbox"/> Microfracture |
| <input type="checkbox"/> Labral debridement | <input type="checkbox"/> Fibrin glue repair |
| <input type="checkbox"/> Labral reconstruction | <input type="checkbox"/> Femoroplasty |

- Capsular repair
- Iliopsoas Release

- Endoscopic Trochanteric Bursa Excision
- Endoscopic Abductor Repair

PHASE I: IMMEDIATE POST-OP (0-3 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> • Minimize pain and inflammation • Protect integrity of repair • Avoid post-operative adhesions • Improve pain-free AROM/PROM within stated parameters • Attain non-analgesic gait with use of device and appropriate weight bearing • Address muscle inhibition • Patient demonstrates independence with initial home exercise program
Weight Bearing	<ul style="list-style-type: none"> • Partial weightbearing 20 lbs, step-to pattern, foot flat gait with crutches
Range of Motion Limitations	<ul style="list-style-type: none"> • Hip Flexion: 0-90 deg (PROM) • Hip Extension: 0 degrees, no motion beyond neutral • Hip Abduction: 0-30 degrees • Hip External Rotation: 0-30 degrees (avoid excessive external rotation) • Hip Internal Rotation: 0-30 degrees
Precautions/Guidelines	<ul style="list-style-type: none"> • No active straight leg raises or flexion-based hip exercises • Avoid ambulation to fatigue or if painful • No active hip flexion post-op days 0-21, hip flexion should be self-assisted for functional mobility • No Gr III-IV hip joint mobilization for until week 8 • No long axis hip distraction for until week 8 for labral repair • No long axis hip distraction until week 12 for labral reconstruction • Avoid pain and pinching in the hip at all times <p>Throughout rehabilitation period every effort should be made to avoid:</p> <ul style="list-style-type: none"> • Hip flexor tendonitis • Synovitis of operative joint • Trochanteric bursitis • Lower back pain or sacroiliac pain
Interventions	<p>Patient Education</p> <ul style="list-style-type: none"> • Activity modification, bed mobility, positioning: <ul style="list-style-type: none"> ○ No crossing of legs ○ Avoid sitting for more than 30 minutes at a time for first 2 weeks and vary position frequently throughout the day. Gradually increase sitting time as tolerated after first 2 weeks. ○ Sit with hip angle less than 90 degrees by sitting on a highchair or sit slightly reclined ○ Prone lying 15 minutes 2-3 times per day to avoid hip flexor contracture ○ Assist operative leg when getting in/out of bed, in/out of car and for all functional mobility ○ Consider obtaining raised toilet seat to avoid hip flexion greater than 90 degrees when sitting on toilet <p><i>Manual Therapy</i></p> <ul style="list-style-type: none"> • Soft tissue mobilization as appropriate for thigh, hip, and lumbar spine musculature. Avoid suture sites until sutures removed and incisions healed • Joint mobilizations to lumbar spine/sacrum to address lumbosacral dysfunction as indicated • Gr I-II hip joint mobilizations for pain modulation as appropriate • Initiate small range hip circumduction and passive IR as indicated below <p><i>Range of motion/Mobility</i></p>

	<ul style="list-style-type: none"> PROM small range hip circumduction at 70° Hip Flexion PROM log rolls to internal rotation/external rotation CPM 2 hours/day <p><i>Gait Training</i></p> <ul style="list-style-type: none"> Gait training with B axillary crutches maintaining indicated weight bearing Stair training with step to pattern, maintaining indicated weight bearing with rail/assistive device <p><i>Modalities</i></p> <ul style="list-style-type: none"> Cryotherapy as needed Electrical stimulation for pain management as needed <p><i>Therapeutic Exercise</i></p> <ul style="list-style-type: none"> Supine Ankle Pumps Supine Quad Set Supine Glute Set Prone Knee Flexion Passive Supine Hip Flexor Stretch (avoid excessive extension) <p><i>Core strengthening</i></p> <ul style="list-style-type: none"> Transversus Abdominis Activation Hooklying Stool rotations hip AAROM ER/IR <p>Hip isometrics</p> <ul style="list-style-type: none"> NO FLEXION Abduction, adduction, ER, IR <p>Cardiovascular Exercise</p> <ul style="list-style-type: none"> Upright Stationary Bike 20 minutes per day, up to 2x/day
Criteria to Progress	<ul style="list-style-type: none"> Minimal pain with ambulation Non-antalgic gait with use of crutches Minimal pain at rest Patient able to perform exercise program without increase in baseline pain Patient compliant with weight bearing, home exercise program, and activity precautions

PHASE II: INTERMEDIATE POST-OP (4-6 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> Progress weight bearing as appropriate per timeline Progress ROM as tolerated per protocol Minimize pain and inflammation Protect integrity of repair Avoid post-operative adhesions Improve pain-free AROM/PROM within stated parameters Attain non-antalgic gait with use of device and appropriate weight bearing Address muscle inhibition Patient demonstrates independence with initial home exercise program
Weight Bearing	<ul style="list-style-type: none"> Gradually increase weight bearing to WBAT pain-free
Range of Motion Limitations	<ul style="list-style-type: none"> Flexion: gradually increase in pain free manner Extension: 0 -10 degrees Abduction: 0-45 degrees External Rotation: 0-45 degrees Internal Rotation: 0-45 degrees
Precautions/Guidelines	<ul style="list-style-type: none"> No active straight leg raises for 8 weeks No Gr III-IV hip joint mobilization for 1st 6 weeks No long axis hip distraction for first 8 weeks for labral repair No long axis hip distraction for first 12 weeks for labral reconstruction Avoid pain and pinching in the hip at all times

<p>Additional Intervention *Continue with Phase I interventions, as indicated</p>	<ul style="list-style-type: none"> • Avoid functional activities that cause hip pain <p><i>Manual Therapy</i></p> <ul style="list-style-type: none"> • Joint mobilizations to lumbar spine/sacrum to address lumbosacral dysfunction as indicated • Gr I-II hip joint mobilizations as appropriate • Scar mobilization to portal scars as appropriate • PROM small range hip circumduction at 70 degrees flexion • PROM log rolls to internal rotation/external rotation • PROM all motions within allowed ROM <p><i>Gait Training</i></p> <ul style="list-style-type: none"> • Increase to weightbearing as tolerated with bilateral axillary crutches and normalize gait pattern. <ul style="list-style-type: none"> ◦ Avoid contralateral pelvic drop. • May begin to wean to single crutch at 1-2 week • Wean off crutches by 6-8 weeks as tolerated <p><i>Modalities</i></p> <ul style="list-style-type: none"> • Cryotherapy as needed • Electrical stimulation for pain management as needed <p><i>Therapeutic Exercise</i></p> <ul style="list-style-type: none"> • Hip rotations on stool IR/ER • Prone B hip IR • Hook-lying Lumbar Rotation (small range) • Hip ABD/ADD Isometrics Hook-lying • Hook-lying Gluteal Set • Standing Knee Flexion • Supine Modified Thomas Stretch (operative leg straight) • Sidelying Piriformis Stretch • Bilateral Bridging • Standing Hip Abduction • Standing Hip Extension to Neutral • Sidelying Clamshell in Neutral • Hip Internal Rotation Prone with Resistance <p><i>Core strengthening</i></p> <ul style="list-style-type: none"> • Quadruped Rocking • Quadruped Hip Extension Knee Slides for Operative Leg w/TrA Activation • Quadruped 'Cat and Camel' Exercise • Quadruped Hip Extension for Operative Leg • Counter Plank • Sub-maximum pain free hip flexion isometrics (week 4) <p><i>Proprioceptive exercise (week 4)</i></p> <ul style="list-style-type: none"> • Balance boards • Single leg balance • BOSU training (bilateral stance) • Airex pads <p><i>Cardiovascular Exercise</i></p> <ul style="list-style-type: none"> • Upright bike up to 20 minutes, 2x per day with seat slightly elevated to minimize excessive hip flexion, no resistance • Elliptical at 3-4 wks. • Treadmill side stepping (week 4)
--	--

	<ul style="list-style-type: none"> • May begin aqua therapy in low end water at week 4 once incision is healed (avoid swimming/treading)
Criteria to Progress	<ul style="list-style-type: none"> • ROM within functional limits (>90% of uninvolved side) • Ascend/descend 8-inch step with good pelvic control • Good pelvic control during single-limb stance • Normalized gait without an assistive device • No joint inflammation, muscular irritation, or pain • Good neuromuscular control and optimal muscle firing patterns

PHASE III: LATE POST-OP (7-12 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> • Performance of exercise program without hip pain • Normalize hip ROM through appropriate ROM progression as outlined • Good activation of hip musculature without evidence of muscle inhibition • Normalized soft tissue of hip and lumbopelvic region • Normal gait without evidence of gait deviations
Weight Bearing	<ul style="list-style-type: none"> • 6-8 weeks post-op: Gradually wean off crutches
Range of Motion Limitations	<ul style="list-style-type: none"> • Continue to increase hip flexion gradually in a pain-free manner (beginning with isometrics) • Increase hip extension, abduction, external rotation, and internal rotation ROM to full as tolerated
Precautions/Guidelines	<ul style="list-style-type: none"> • No extreme combined ROM (e.g. flexion/IR, flexion/ER) • No plyometrics • No running • No squatting below 90 degrees • Avoid painful ROM • No pivoting on operative leg • Avoid extreme combined hip ROM • Avoid symptom provocation during ambulation, ADLs, or therapeutic exercise and avoid post-activity soreness • Avoid pinching in operative hip with range of motion exercises
Additional Interventions <i>*Continue with Phase I-II interventions, as indicated</i>	<p><i>Manual Therapy</i></p> <ul style="list-style-type: none"> • Gr III-IV hip joint mobilization as needed to address joint hypomobility • Long axis hip distraction if needed beginning at 8 weeks for labral repair • No long axis hip distraction for first 12 weeks for labral reconstruction <p><i>Gait Training</i></p> <ul style="list-style-type: none"> • Normalize gait without device by 6-8 weeks • If patient has pain with ambulation continue to use 1 crutch and wean as tolerated <p><i>Modalities</i></p> <ul style="list-style-type: none"> • Cryotherapy as needed • Electrical stimulation for pain management as needed. <p><i>Therapeutic Exercise</i></p> <ul style="list-style-type: none"> • Submaximal hip flexor isometrics • Sidelying Hip Abduction • Partial Range Squats (gradually increase to 90 degree squats) • Prone Hip Extension • Single Leg Forward Weight Shifts (progressing to Romanian dead lift) • Lateral Band Walk • Backwards Monster Walk with Band • Banded Hip Clamshell • Standing IT Band Stretch

	<ul style="list-style-type: none"> • Leg press (avoiding deep flexion) • Hip hiking on Stairmaster or off step <p><i>Core Strengthening</i></p> <ul style="list-style-type: none"> • Paloff Press • Bridge with Alternating Leg Extension • Side Plank- modified (knees/forearm) • Modified Plank (knees/forearms) • Quadruped Alternating Leg Extension (progress to opposite arm/leg as tolerated) <p><i>Proprioceptive exercise (avoid excessive torsion in hip until wk. 8)</i></p> <ul style="list-style-type: none"> • Single Leg Balance with Clock Taps • Single Leg Balance with Hip ABD and Band Resistance • Single Leg Balance with Hip Ext and Band Resistance <p><i>Cardiovascular Exercise</i></p> <ul style="list-style-type: none"> • Upright stationary bicycle: gradually increase time and resistance as tolerated • Elliptical training: pedaling forward and backward if pain-free, gradually increase time and resistance as tolerated • Swimming: initiate flutter kick as tolerated, avoid frog kicking • No treadmill walking/running
Criteria to Progress	<ul style="list-style-type: none"> • ROM within limits pain-free • Alternate Ascend/Descend 8-inch step with good pelvic control no UE support • Good pelvic control during single-limb stance and dynamic balance • Normalized gait pain-free without an assistive device • No Pain at rest, ADL/IADL nor walking • Strength of operative hip 75% of contralateral hip • No joint inflammation, muscular irritation, or pain • Good neuromuscular control and optimal muscle firing patterns

PHASE IV: ADVANCED REHABILITATION (12-16 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> • Independent home exercise program • Optimize ROM • $\geq 4/5$ LE strength, $\geq 4/5$ trunk strength • Improved dynamic balance • Pain-free ADL • Pain-free hip flexion with ADLs and functional mobility • May begin treadmill running at 16 weeks if cleared by surgeon • If full hip ROM still not attained, continue to progress as tolerated
Precautions	<ul style="list-style-type: none"> • No extreme combined ROM (e.g. flexion/IR, flexion/ER) • No plyometrics • No squatting below 90 degrees • Avoid painful ROM • Avoid extreme combined hip ROM • No symptom provocation during ambulation, ADLs, or therapeutic exercise • Avoid pinching in operative hip with range of motion exercises
Additional Interventions <i>*Continue with Phase I-III interventions, as indicated</i>	<p><i>Manual Therapy</i></p> <ul style="list-style-type: none"> • Long axis hip distraction if needed <p><i>Modalities</i></p> <ul style="list-style-type: none"> • Cryotherapy as needed • Electrical stimulation for pain management as needed <p><i>Therapeutic Exercise</i></p> <ul style="list-style-type: none"> • Progressive lower extremity and core exercises by increasing challenge and resistance

	<ul style="list-style-type: none"> Advanced balance exercises as appropriate for sport or desired recreation Sport specific plyometrics and agility exercises as appropriate Progress core strengthening as deemed appropriate by therapist <p><i>Cardiovascular Exercise</i></p> <ul style="list-style-type: none"> Treadmill running (if cleared) to begin at 14-16 weeks, see return to run program
Criteria to Progress	<ul style="list-style-type: none"> Y Balance Test Limb symmetry index 80% of uninjured side Strength of operative hip 90% of uninjured side Perform progressed exercise program without pain No joint inflammation, muscular irritation, or pain

PHASE V: RETURN TO SPORT (>16 WEEKS AFTER SURGERY)

Rehabilitation Goals	<ul style="list-style-type: none"> Progress to sport specific training without pain Progress to jogging pain free when cleared by surgeon Independent home exercise program Optimize ROM 5/5 LE strength, \geq4/5 trunk strength Normal Muscle Length of B LE Good, dynamic unilateral balance of operative extremity Pain-free with all activities
Precautions/Guidelines	<ul style="list-style-type: none"> Avoid pain in hip joint with functional activities or exercises If post-exercise joint pain or limping occurs, activity level should be modified Avoid joint inflammation Focus on quality of movement and exercise
Additional Interventions <i>*Continue with Phase I-IV interventions, as indicated</i>	<p><i>Modalities</i></p> <ul style="list-style-type: none"> Cryotherapy as needed Electrical stimulation for pain management as needed <p><i>Therapeutic Exercise</i></p> <ul style="list-style-type: none"> Progress strength, proprioception, dynamic balance, agility, and power to address sport specific demands. Sport specific retraining as tolerated. <p><i>Cardiovascular Exercise</i></p> <ul style="list-style-type: none"> Jogging: may initiate walk/jog interval return to run program once appropriate criteria have been met and has been cleared by surgeon at week 16
Criteria to Progress	<ul style="list-style-type: none"> Cross over triple hop for distance 90% of uninjured side Y Balance Test Limb symmetry index 80% of uninjured side Quad/hamstrings/gluteal index \geq 90% HHD testing Patient able to jog 30 minutes Patient able to perform sport specific drills without pain Good neuromuscular control and optimal muscle firing patterns <p>Outcome Measures:</p> <ul style="list-style-type: none"> Hip Outcome Score (HOS) <ul style="list-style-type: none"> Lower Extremity Functional Scale (LEFS) may also be used as an alternative

Revised 3/2023

Contact	Please email MGHSportsPhysicalTherapy@partners.org with questions specific to this protocol
----------------	--

References:

- Dippmann, C., Thorborg, K., Kraemer, O., Winge, S., Palm, H., & Hölmich, P. (2014). Hip arthroscopy with labral repair for femoroacetabular impingement: short-term outcomes. *Knee Surgery, Sports Traumatology, Arthroscopy*, 22(4), 744-749.
- Kuhns, B. D., Weber, A. E., Batko, B., Nho, S. J., & Stegemann, C. (2017). A four-phase physical therapy regimen for returning athletes to sport following hip arthroscopy for femoroacetabular impingement with routine capsular closure. *International journal of sports physical therapy*, 12(4), 683.

3. Heerey, J., Risberg, M. A., Magnus, J., Moksnes, H., Ødegaard, T., Crossley, K., & Kemp, J. L. (2018). Impairment-based rehabilitation following hip arthroscopy: postoperative protocol for the HIP ARthroscopy International Randomized Controlled Trial. *Journal of orthopaedic & sports physical therapy*, 48(4), 336-342.
4. Willimon, S. C., Briggs, K. K., & Philippon, M. J. (2014). Intra-articular adhesions following hip arthroscopy: a risk factor analysis. *Knee Surgery, Sports Traumatology, Arthroscopy*, 22(4), 822-825.
5. Enseki, K. R., & Kohlrieser, D. (2014). Rehabilitation following hip arthroscopy: an evolving process. *International journal of sports physical therapy*, 9(6), 765.
6. Enseki, K. R., Martin, R., & Kelly, B. T. (2010). Rehabilitation after arthroscopic decompression for femoroacetabular impingement. *Clinics in sports medicine*, 29(2), 247-255.
7. Philippon, Marc J. MD; Yen, Yi-Meng MD, PhD; Briggs, Karen K. MBA, MPH; Koppersmith, David A. BS; Maxwell, Richard Brian BS. (2008) Early Outcomes After Hip Arthroscopy for Femoroacetabular Impingement in the Athletic Adolescent Patient: A Preliminary Report. *Journal of Pediatric Orthopedics*, 28(7), 705-710.
8. Pathy, Rubinia; Sink, Ernest L.b. (2016). Femoroacetabular impingement in children and adolescents. *Current Opinion in Pediatrics*, 28(1), 68-78.
9. Ayeni OR, Karlsson J, et al. (2021) Osteochondroplasty and Labral Repair for the Treatment of Young Adults With Femoroacetabular Impingement: A Randomized Controlled Trial. *The American Journal of Sports Medicine*, 49(1), 25-34.
10. Smith, David V.; Bernhardt, David T.. Hip Injuries in Young Athletes. (2010). *Current Sports Medicine Reports*, 9(5), 278-283.
11. Emily Arciero, Rafael Kakazu, Patrick Garvin, Allison E. Crepeau, Katherine Coyner. (2022). Favorable Patient-Reported Outcomes and High Return to Sport Rates Following Hip Arthroscopy in Adolescent Athletes: A Systematic Review, *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, 38(9), 2730-2740.
12. Damian R Griffin, Edward J Dickenson, et. al. (2018). Hip arthroscopy versus best conservative care for the treatment of femoroacetabular impingement syndrome (UK FASHION): a multicenter randomized controlled trial, *The Lancet*, 391(10136), 2225-2235.